

## On Gödel's incompleteness theorems

S. Kalimuthu

2/394, Kanjampatti P.O., Pollachi via, Tamilnadu 642003, India

Email: [sona.sonasona.sona7@gmail.com](mailto:sona.sonasona.sona7@gmail.com)

### Abstract

In this short communication, the mathematical variation of the Liar's paradox of the Gödelian incompleteness theorem was proved.

**MSC:** 51 M 04

**Key Words:** Euclidean postulates, Gödelian incompleteness theorem

### 1 Introduction

In 1931, an young Austrian mathematician Kurt Gödel published a paper in mathematical logic. In this ground breaking paper, he has proved two propositions. Gödel's findings are called Gödel's incompleteness theorems. This work was a masterstroke for Hilbert's second theorem. Gödel's theorems are given below:

**Theorem 1:** In any logical system one can construct statements that are neither true nor false (mathematical variations of the liar's paradox).


**Theorem 2:** Therefore no consistent system can be used to prove its own consistency. No proof can be proof of itself. [<http://milesmathis.com/godel.htm>]

In this short work, we are going to establish the first theorem.

### 2 Construction

Draw triangles ABC and DBC as shown in figure 1. On AB, choose a point E. Join C and E. Join E and D meeting BC at R. Since points E and D lie on the opposite sides of BC, ED can meet BC. Please note that Euclid uses this principle. [ Elements I, prop.10 ] Similarly join A

and D contacting EC at S and RC at O. Small letters denote the angle sums of triangles and quadrilateral ORES. Also, let that  $a, b, c, d, e, f, g, h, i, j, k, l, t$  and  $w$  respectively refer to the sum of the interior angles of triangles and quadrilaterals ACD, ACO, CDS, AED, SED, AERO, BDE, AEC, EBC, ERC, BOSE, BCD, CDR and BOD.



**Euclidean figure 1**

### 3 Results

The angles BRO, ROC, ERD, ASO, SOD, ESC and AEB are all straight angles and so their measures are all equal to 180 degrees. Let  $v$  be the value of this 180 degrees. (1)

Applying (1),  $x + y + z = 2v + a$  (2)

$$x + y = v + b \quad (3)$$

$$y + z = v + c \quad (4)$$

$$m + n + p = 3v + d \quad (5)$$

$$m + n = 2v + e \tag{6}$$

$$p + n = v + f \tag{7}$$

$$q + r = v + g \tag{8}$$

$$x + p = v + h \tag{9}$$

$$y + n + q = 3v + i \tag{10}$$

$$y + n = 2v + j \tag{11}$$

$$n + q = v + k \tag{12}$$

$$z + m + r = 2v + l \tag{13}$$

$$z + m = v + t \tag{14}$$

$$m + r = v + w \tag{15}$$

Adding (2) to (15) we have,  $3x + 5y + 4z + 5m + 6n + 3p + 3q + 3r =$

$$22v + a + b + c + d + e + f + g + h + i + j + k + l + t + w$$

Putting (4) , (6) , (8) and (9) in LHS,  $2h + 2g + 3c + 4e + y + n =$

$$2v + a + b + d + f + i + j + k + l + t + w$$

Applying (11) in LHS ,  $2h + 2g + 3c + 4e = a + b + d + f + i + k + l + t + w \tag{16}$

$(4) + (6) = (11) + (14) = y + z + m + n = 3v + c + e = 3v + j + t, \text{ i.e } c + e = j + t \tag{17}$

Applying (17) in (16) we obtain that,  $2g + 2h + 2c + 3e + j =$

$$a + b + d + f + i + k + l + w$$

$$2v + a = x + y + z \tag{2}$$

$$v + b = x + y \tag{3}$$

$$3v + d = m + n + p \tag{5}$$

$$v + f = p + n \tag{7}$$

$$3v + i = y + n + q \tag{10}$$

$$v + k = n + q \tag{12}$$

$$2v + l = z + m + r \quad (13)$$

$$v + w = n + r \quad (15)$$

$$y + n = 2v + j \quad (11)$$

$$m + n = 2v + e \quad (6)$$

Adding the above eleven relations,  $10v + 2g + 2h + 2c + 3e =$

$$2x + 2y + 2z + m + 3n + 2q + 2p + 2r$$

Assuming (4), (6), (8) and (9) in RHS,  $2v + 2e = 2n$ , i. e.  $n = v + e$  (18)

Applying (18) in (6) we obtain that,  $m = v$  (19)

Comparing (1) and  $m$  we get that the sum of the interior angles of triangle

ORD is equal to two right angles (20)

Needless to say, (20) proves Euclid's fifth postulate. [1 – 4] (21)

#### 4 Discussions

It is well known that Beltrami, Klein, Cayley, Poincare and others have shown that it is impossible to deduce Euclid V from Euclid I to IV. But the author's equation (21) proves the fifth Euclidean postulate. A brief analysis of these two theorems shows that it established Gödel's first incompleteness theorem mentioned above.

#### References

- [1] Kalimuthu, Pretty algebra , Marsland Press , New York , 2009 , 7(6) , pp 86 to 89
- [2] Kalimuthu, Beautiful geometry, Marsland Press , New York , 2009 , 7(7) , pp 88 to 89
- [3] Kalimuthu, For the origin of new geometry, Applied Mathematics Letter, 2010, 23(1), 1451-1453
- [4] Kalimuthu, On applied algebra, Advances in Algebra, 2009, 2(1), 41-43